

## AUTHOR INDEX

- Aitchison S. See Blaylock JF.
- Akin T, Aykan U, Yildiz TF, Karadayi K, Bilge AH. Intraocular lens tilt and decentration after phacoemulsification surgery in patients with and without primary open angle glaucoma. 24(9):865. [Letter]
- Albè E. See Vinciguerra P.
- Albelda AE. See Palmer AM.
- Alfonso JF, Palacios A, Montés-Micó R. Myopic phakic STAAR collamer posterior chamber intraocular lenses for keratoconus. 24(9):867-874.
- Alhasso D. See Elsheikh A.
- Alió JL, Piñero DP. Very high-frequency digital ultrasound measurement of the LASIK flap thickness profile using the IntraLase femtosecond laser and M2 and Carriazo-Pendular microkeratomes. 24(1):12-23.
- Alió JL. See Illueca C.
- Alió JL. See Javaloy J.
- Alió JL. See Patel S.
- Alió JL. See Pinelli R.
- Alió JL. See Uceda-Montanes A.
- Alonso RS. See Fontes BM.
- Ambrósio R Jr, Tervo T, Wilson SE. LASIK-associated dry eye and neurotrophic epitheliopathy: pathophysiology and strategies for prevention and treatment. 24(4):396-407. [Review]
- Ambrósio R Jr. See Fontes BM.
- Anderson DM. See McCauley MB.
- Ando W. See Kamiya K.
- Applegate RA, Krueger RR. Introduction to the Proceedings of the 9th International Congress of Wavefront and Presbyopic Refractive Corrections. 24(9):963-964. [Proceedings]
- Arba-Mosquera S. See de Ortueta D.
- Archer TJ. See Reinstein DZ.
- Arné JL. See Couillet J.
- Arranz-Marquez E. See Gil-Cazorla R.
- Asato Y. See Kamiya K.
- Aslanides IM. See Mearza AA.
- Auffarth G. See Scheuerle AF.
- Awwad ST, Warmerdam D, Bowman RW, Dwarakanathan S, Cavanagh HD, McCulley JP. Contrast sensitivity and higher order aberrations in eyes implanted with AcrySof IQ SN60WF and AcrySof SN60AT intraocular lenses. 24(6):619-625.
- Aykan U. See Akin T.
- Baatz H. See de Ortueta D.
- Bacchi C. See Pinelli R.
- Bains HS. See Dougherty PJ.
- Balestrazzi A. See Casprini F.
- Bamba S. See Ramos-Esteban JC.
- Barboni P. See Savini G.
- Barequet IS, Hirsh A, Levinger S. Effect of thin femtosecond LASIK flaps on corneal sensitivity and tear function. 24(9):897-902.
- Barrera C. See Javaloy J.
- Barreto J Jr. See Netto MV.
- Battle JF. See Peyman GA.
- Bayer A. See Erdem U.
- Bechara S. See Netto MV.
- Behrens A, Stark WJ, Pratzter KA, McDonnell PJ. Dynamics of small-incision clear cornea wounds after phacoemulsification surgery using optical coherence tomography in the early postoperative period. 24(1):46-49. [Report]
- Belin MW. See Khachikian SS.
- Belmonte C. See Tuisku IS.
- Bernitsky DA. See Sanders DR.
- Bharadwaj SR. See Schor CM.
- Bilge AH. See Akin T.
- Bilgihan K. See Hondur A.
- Binder PS. See Warden L.
- Blaylock JF, Si Z, Aitchison S, Prescott C. Visual function and change in quality of life after bilateral refractive lens exchange with the ReSTOR multifocal intraocular lens. 24(3):265-273.
- Boccuzzi D. See Cennamo G.
- Boghossian AJ. See Stahl JE.
- Bonilha VL. See de Medeiros FW.
- Borrelli M, Irregolare C. Calculating the keratometric refractive index. 24(1):10. [Letter]
- Bottós KM, Dreyfuss JL, Regatieri CVS, Lima-Filho AAS, Schor P, Nader HB, Chamon W. Immunofluorescence confocal microscopy of porcine corneas following collagen cross-linking treatment with riboflavin and ultraviolet A. 24(7):S715-S719. [Proceedings]
- Bouzoukis D. See Kymionis GD.
- Bowman RW. See Awwad ST.
- Bowman RW. See Petroll WM.
- Boxer Wachler B. Metrics for keratoconus. 24(5):460. [Letter]
- Braunstein RE. See Paik DC.
- Bricola G. See Iester M.
- Brown M. See Elsheikh A.
- Bueeler M. See Mrochen M.
- Bunce C. See Wang JC.
- Cabrera G. See Peyman GA
- Cadarsó L, Iglesias A, Ollero A, Pita B, Montés-Micó R. Postoperative optical aberrations in eyes implanted with AcrySof spherical and aspheric intraocular lenses. 24(8):811-816.
- Cakir H. See Utine CA.
- Calabria G. See Iester M.
- Caldwell M, Reilly C. Effects of topical nepafenac on corneal epithelial healing time and postoperative pain after PRK: a bilateral, prospective, randomized, masked trial. 24(4):377-382.
- Calossi A. Mathematical properties of asphericity: a method to calculate with asphericities. 24(2):121. [Reply to Letter]
- Camellin M, Wyler D. Epi-LASIK versus epi-LASEK. 24(1):S57-S63. [Proceedings]
- Camellin M. What about LASEK? 24(5):462. [Letter]
- Campanelli M. See Elsheikh A.
- Campbell CE. Wavefront measurements of diffractive and refractive multifocal intraocular lenses in an artificial eye. 24(3):308-311. [Report]
- Campos M. See Wallau AD.
- Caporossi A. See Casprini F.
- Caporossi T. See Casprini F.
- Casprini F, Balestrazzi A, Tosi GM, Lazzarotto M, Malandrini A, Lepri F, Martone G, Caporossi T, Caporossi A. Optical aberrations in pseudophakic eyes after 2.5-mm Nd:YAG laser capsulotomy for posterior capsule opacification. 24(7):702-706.
- Cavanagh HD. See Awwad ST.
- Cavanagh HD. See Petroll WM.

- Celik L. See Gunenc U.
- Cennamo G, Intravaja A, Boccuzzi D, Marotta G, Cennamo G. Treatment of keratoconus by topography-guided customized photorefractive keratectomy: two-year follow-up study. 24(2):145-149.
- Cennamo G. See Cennamo G.
- Cerviño A, Hosking SL, Montés-Micó R. Comparison of higher order aberrations measured by NIDEK OPD-Scan dynamic skiascopy and Zeiss WASCAs Hartmann-Shack aberrometers. 24(8):790-796.
- Céspedes MC. See Palmer AM.
- Chamacham T. See Ghaffariyeh A.
- Chamon W. See Bottós KM.
- Chan TK. See Lim MCC.
- Chang JSM. Complications of sub-Bowman's keratomileusis with a femtosecond laser in 3009 eyes. 24(1):S97-S101. [Proceedings]
- Chang SW, Chou SF, Chuang JL. Mechanical corneal epithelium scraping and ethanol treatment up-regulate cytokine gene expression differently in rabbit cornea. 24(2):150-159.
- Charisis S. See Kymionis GD.
- Charman WN, Montés-Micó R, Radhakrishnan H. Problems in the measurement of wavefront aberration for eyes implanted with diffractive bifocal and multifocal intraocular lenses. 24(3):280-286.
- Chaurasia SS. See de Medeiros FW.
- Chavan R. See Savant V.
- Chayet A. See Waring GO III.
- Chen H. See Yu J.
- Chen WL. See Lin JM.
- Cheng ACK, Ho T, Lau S, Wong AL, Leung C, Lam DSC. Measurement of LASIK flap thickness with anterior segment optical coherence tomography. 24(9):879-884.
- Cheng ACK, Ho T, Rao SK, Lau S, Lam DSC. Posterior corneal curvature measurements with peripheral fitting zones before and after myopic LASIK using Orbscan II. 24(8):807-810.
- Cheng ACK, Rao SK, Lau S, Leung CKS, Lam DSC. Central corneal thickness measurements by ultrasound, Orbscan II, and Visante OCT after LASIK for myopia. 24(4):361-365.
- Cheng ACK, Rao SSK, Lau S, Wong A, Lam DSC. Comparison of techniques for corneal power assessment after myopic LASIK without the use of preoperative data. 24(5):539-543.
- Cheng ACK, Rao SSK, Lau S, Wong A, Lam DSC. Corneal power after myopic LASIK. 24(8):770. [Reply to Letter]
- Cheng ACK. See Srivannaboon S.
- Cheng Z. See Liu Z.
- Cheng ZY, He JC, Zhou XT, Chu RY. Effect of flap thickness on higher order wavefront aberrations induced by LASIK: a bilateral study. 24(5):524-529.
- Chiang CC. See Lin JM.
- Choi CY, Kim JY, Kim MJ, Tchah H. Transmission electron microscopy study of corneal epithelial flaps following removal using mechanical scraping, alcohol, and epikeratome techniques. 24(7):667-670.
- Choi J, Schwiegerling J. Optical performance measurement and night driving simulation of ReSTOR, ReZoom, and Tecnis multifocal intraocular lenses in a model eye. 24(3):218-222.
- Choi J. See Schwiegerling J.
- Chokshi A. See Khalil M.
- Chou SF. See Chang SW.
- Chu RY. See Cheng ZY.
- Chuang JL. See Chang SW.
- Chung SA, Kim EK, Ryu IKH, Kim JK, Lee HK. Effectiveness of cultured human keratinocyte onlays on epithelial healing and clinical outcome after photorefractive keratectomy. 24(8):826-832.
- Ciolino JB. See Khachikian SS.
- Coleman DJ. See Reinstein DZ.
- Coral SA. See Ghanem VC.
- Corcostegui I. See Maldonado MJ.
- Coullet J, Fournié P, Malecaze F, Arné JL. Inadequate results for microkeratome-assisted additive stromal keratoplasty for management of keratoconus. 24(2):166-172.
- Dawson DG, Grossniklaus HE, McCarey BE, Edelhauser HF. Biomechanical and wound healing characteristics of corneas after excimer laser keratorefractive surgery: is there a difference between advanced surface ablation and sub-Bowman's keratomileusis? 24(1):S90-S96. [Proceedings]
- Dawson DG. See Randleman JB.
- de Benito-Llopis L, Teus MA, Sánchez-Pina JM. Comparison between LASEK with mitomycin C and LASIK for the correction of myopia of -7.00 to -13.75 D. 24(5):516-523.
- De Feo F. See Iester M.
- de Freitas D. See El Dib RP.
- de Freitas D. See Vieira AC.
- de Medeiros FW, Mohan RR, Suto C, Sinha S, Bonilha VL, Chaurasia SS, Wilson SE. Haze development after photorefractive keratectomy: mechanical vs ethanol epithelial removal in rabbits. 24(9):923-927.
- de Ortueta D. Planar flaps with the Carriazo-Pendular microkeratome. 24(4):322. [Letter]
- de Ortueta D, Arba-Mosquera S, Baatz H. Topographic changes after hyperopic LASIK with the SCHWIND ESIRIS laser platform. 24(2):137-144.
- de Ortueta D, Mosquera SA. Mathematical properties of asphericity: a method to calculate with asphericities. 24(2):119-121. [Letter]
- de Sanctis U, Mutani B, Grignolo FM. Long-term endothelial cell loss after traumatic dislocation and repositioning of artisan phakic IOL. 24(5):546-548.
- de Souza DC. See Ghanem RC.
- Di Iorio E. See Vinciguerra P.
- Diakonis VF. See Kymionis GD.
- Donitzky C. See Mrochen M.
- Donnelly W III. The advanced human eye model (AHM): a personal binocular eye modeling system inclusive of refraction, diffraction, and scatter. 24(9):976-983. [Proceedings]
- Dougherty PJ, Bains HS. A retrospective comparison of LASIK outcomes for myopia and myopic astigmatism with conventional NIDEK versus wavefront-guided VISX and Alcon platforms. 24(9):891-896.
- Drake A. See Srinivasan S.
- Dreher AW. See Warden L.
- Dreyfuss JL. See Bottós KM.
- Durrie DS, Slade SG, Marshall J. Wavefront-guided excimer laser ablation using photorefractive keratectomy and sub-Bowman's keratomileusis: a contralateral eye study. 24(1):S77-S84. [Proceedings]
- Durrie DS. See Eser I.
- Durrie DS. See Stahl JE.
- Durrie DS. See Waring GO IV.
- Swarakanathan S. See Awwad ST.
- Eckhardt HB. See Hütz WW.
- Edelhauser HF. See Dawson DG.
- Edelhauser HF. See Randleman JB.
- Egemenoglu A. See Utine CA.
- Ehrenhaus MP. See Ursea R.

- El Dib RP, de Freitas D. A systematic review of Ferrara's ring in the treatment of keratoconus. 24(9):865-866. [Letter]
- Elsheikh A, Brown M, Alhasso D, Rama P, Campanelli M, Garway-Heath D. Experimental assessment of corneal anisotropy. 24(2):178-187. [Biomechanics]
- Erdem U, Muftuoglu O, Gundogan FC, Sobaci G, Bayer A. Pupil center shift relative to the coaxially sighted corneal light reflex under natural and pharmacologically dilated conditions. 24(5):530-538.
- Ertan A, Karacal H. Factors influencing flap and INTACS decentration after femtosecond laser application in normal and keratoconic eyes. 24(8):797-801.
- Ertan A, Ozkiloglu E. Effect of age on outcomes in patients with keratoconus treated by Intacs using a femtosecond laser. 24(7):690-695.
- Ertan A. See Kamburoglu G.
- Eser I, Durrie DS, Schwendeman F, Stahl JE. Association between ocular dominance and refraction. 24(7):685-689.
- Espinosa J. See Illueca C.
- Fahle M. See Kaymak H.
- Faiña PG. See Palmer AM.
- Fam HB, Lim KL. A comparative analysis of intraocular lens power calculation methods after myopic excimer laser surgery. 24(4):355-360.
- Fam HB. See Srivannaboon S.
- Fant B. See Waring GO III.
- Feinbaum C. See Patel S.
- Féliz R. See Peyman GA.
- Fischer J. See Waring GO III.
- Fontes BM, Ambrósio R Jr, Alonso RS, Jardim D, Velarde GC, Nosé W. Corneal biomechanical metrics in eyes with refraction of -19.00 to +9.00 D in healthy Brazilian patients. 24(9):941-945. [Biomechanics]
- Fournié P. See Couillet J.
- Fromm M. See Schumacher S.
- Fujisawa T. See Kamiya K.
- García-Layana A. See Maldonado MJ.
- Garway-Heath D. See Elsheikh A.
- Gerten G. See Schumacher S.
- Ghaffari R. See Hashemi H.
- Ghaffariyeh A, Chamacham T. Tricyclic antidepressants: potential therapeutic alternatives for treatment of dry eye symptoms after LASIK. 24(8):770-771. [Letter]
- Ghanem EA. See Ghanem RC.
- Ghanem RC, Ghanem VC, de Souza DC, Kara-Jose N, Ghanem EA. Customized topography-guided photorefractive keratectomy with the MEL-70 platform and mitomycin C to correct hyperopia after radial keratotomy. 24(9):911-922.
- Ghanem RC. See Ghanem VC.
- Ghanem VC, Kara-José N, Ghanem RC, Coral SA. Photorefractive keratectomy and butterfly laser epithelial keratomileusis: a prospective, contralateral study. 24(7):671-684.
- Ghanem VC, Souza GC, Souza DC, Vieste JMZ, Weber SLP, Kara-José N. PRK and butterfly LASEK: prospective, randomized, contralateral eye comparison of epithelial healing and ocular discomfort. 24(6):591-599.
- Ghanem VC. See Ghanem RC.
- Gil-Cazorla R, Teus MA, Arranz-Márquez E. Comparison of silicone and non-silicone hydrogel soft contact lenses used as a bandage after LASEK. 24(2):199-203. [Report]
- Gil-Cazorla R, Teus MA, Arranz-Marquez E, Marina-Verde C. Phakic refractive lens (Medennium) for correction of +4.00 up +6.00 diopters: 1-year follow-up. 24(4):350-354.
- Ginis H. See Kymionis GD.
- Gobbe M. See Reinstein DZ.
- Goes FJ. Refractive lens exchange with the diffractive multifocal Tecnis ZM900 intraocular lens. 24(3):243-250.
- Goes FJ. Visual results following implantation of a refractive multifocal IOL in one eye and a diffractive multifocal IOL in the contralateral eye. 24(3):300-305. [New Technology]
- Goldman DA. See Ide T.
- Grignolo FM. See de Sanctis U.
- Grolmus R. See Hütz WW.
- Grossniklaus HE. See Dawson DG.
- Grossniklaus HE. See Randleman JB.
- Gundogan FC. See Erdem U.
- Gunenc U, Celik L. Long-term experience with mixing and matching refractive Array and diffractive CeeOn multifocal intraocular lenses. 24(3):233-242.
- Halbhuber KJ. See Wang BG.
- Hamilton DR. See Samimi D.
- Hara S. See Yoshida Y.
- Harton PJ Jr. See Sanders DR.
- Hasanreisoglu B. See Hondur A.
- Hashemi H, Ghaffari R, Mohammadi M, Moghimi S, Miraftaab M. Microbial keratitis after INTACS implantation with loose suture. 24(5):551-552.
- He JC. See Cheng ZY.
- Henriquez MA. See Izquierdo L Jr.
- Herekar S. See Rocha KM.
- Herzig S. See Srinivasan S.
- Hirsh A. See Barequet IS.
- Ho T. See Cheng ACK.
- Hodge C. See Sutton G.
- Hoffer KJ. See Savini G.
- Holland SP. See Lin DTC.
- Holopainen JM. See Neira-Zalentein W.
- Hondur A, Bilgihan K, Hasanreisoglu B. A prospective bilateral comparison of epi-LASIK and LASEK for myopia. 24(9):928-934.
- Hosking SL. See Cerviño A.
- Hütz WW, Eckhardt HB, Röhrig B, Grolmus R. Intermediate vision and reading speed with Array, Tecnis, and ReSTOR intraocular lenses. 24(3):251-256.
- Ichikawa K. See Yoshida Y.
- Ide T, Kymionis GD, Goldman DA, Yoo SH, O'Brien TP. Subconjunctival gas bubble formation during LASIK flap creation using femtosecond laser. 24(8):850-851. [Report]
- Iester M, De Feo F, Bricola G, Papadia M, Venturino G, Traverso CE, Calabria G. Retinal nerve fiber layer measurements before and after photorefractive keratectomy. 24(6):639-644.
- Iglesias A. See Cadarso L.
- Iida Y, Shimizu K, Ito M, Suzuki M. Influence of age on ocular wavefront aberration changes with accommodation. 24(7):696-701.
- Ilango B. See Savant V.
- Illueca C, Alió JL, Mas D, Ortiz D, Pérez J, Espinosa J, Sala E. Pseudoaccommodation and visual acuity with Technovision presbyLASIK and a theoretical simulated Array® multifocal intraocular lens. 24(4):344-349.

- Intravaja A. See Cennamo G.
- Irregolare C. See Borrelli M.
- Iseli HP, Spoerl E, Wiedemann P, Krueger RR, Seiler T. Efficacy and safety of blue-light scleral cross-linking. 24(7):S752-S755. [Proceedings]
- Ito M. See Iida Y.
- Izquierdo L Jr, Henriquez MA, Zakrzewski PA. Detection of an abnormally thick LASIK flap with anterior segment OCT imaging prior to planned LASIK retreatment surgery. 24(2):197-199. [Report]
- Jampaulo M, Maloney RK. Lack of progression of ectasia seven years after LASIK in a highly myopic keratoconic eye. 24(7):707-709. [Report]
- Jankov MR. See Stojanovic A.
- Jardim D. See Fontes BM.
- Javaloy J, Barrera C, Muñoz G, Pérez-Santonja JJ, Vidal MT, Alió JL. Spontaneous bilateral, recurrent, late-onset diffuse lamellar keratitis after LASIK in a patient with Cogan's syndrome. 24(5):548-550. [Report]
- Jendritza BB, Knorz MC, Morton S. Wavefront-guided excimer laser vision correction after multifocal IOL implantation. 24(3):274-279.
- Jiang H. See Liu Z.
- Johnson AJ. See McCauley MB.
- Jonas JB. See Vossmerbaeumer U.
- Joo CK. See Jung SW.
- Joo CK. See Kim H.
- Jung SW, Kim MJ, Park SH, Joo CK. Multifocal corneal ablation for hyperopic presbyopes. 24(9):903-910.
- Kamburoglu G, Ertan A. Intacs implantation with sequential collagen cross-linking treatment in postoperative LASIK ectasia. 24(7):S726-S729. [Proceedings]
- Kamiya K, Shimizu K, Ando W, Asato Y, Fujisawa T. Phakic toric implantable collamer lens implantation for the correction of high myopic astigmatism in eyes with keratoconus. 24(8):840-842. [Report]
- Kanellopoulos AJ. See Krueger RR.
- Kara-José N. See Ghanem RC.
- Kara-José N. See Ghanem VC.
- Kara-José N. See Netto MV.
- Karacal H. See Ertan A.
- Karadayi K. See Akin T.
- Kaymak H, Fahle M, Ott G, Mester U. Intraindividual comparison of the effect of training on visual performance with ReSTOR and Tecnis diffractive multifocal IOLs. 24(3):287-293.
- Kaymak H. See Mester U.
- Kezirian GM, Moore CR, Stonecipher KG, SurgiVision® Consultants Inc WaveLight Investigator Group. Four-year postoperative results of the US ALLEGRETTO WAVE clinical trial for the treatment of hyperopia. 24(4):S431-S438.
- Kezirian GM. See Stonecipher KG.
- Khachikian SS, Belin MW, Ciolino JB. Intrasubject corneal thickness asymmetry. 24(6):606-609.
- Khalil M, Chokshi A, Ltkany R, Speaker MG, Yu G. Prospective evaluation of intraocular lens calculation after myopic refractive surgery. 24(1):33-38.
- Kim EK. See Chung SA.
- Kim EK. See Lee JH.
- Kim EK. See Kim T.
- Kim H, Joo CK. Measure of keratoconus progression using Orbscan II. 24(6):600-605.
- Kim HI. See Park SH.
- Kim JK. See Chung SA.
- Kim JY. See Choi CY.
- Kim MJ. See Choi CY.
- Kim MJ. See Jung SW.
- Kim SW. See Kim T.
- Kim SY. See Park SH.
- Kim T, Kim T, Kim SW, Kim EK. Comparison of corneal deposits after LASIK and PRK in eyes with granular corneal dystrophy type II. 24(4):392-395.
- Kim T. See Kim T.
- Kim WC. See Lee JH.
- Kimura I. See Yamaguchi T.
- Kirwan C, O'Keefe M. Measurement of intraocular pressure in LASIK and LASEK patients using the Reichert Ocular Response Analyzer and Goldmann applanation tonometry. 24(4):366-370.
- Klimava A. See Reznik J.
- Knorz MC, Vossmerbaeumer U. Comparison of flap adhesion strength using the Amadeus microkeratome and the IntraLase iFS femtosecond laser in rabbits. 24(9):875-878.
- Knorz MC. Multifocal intraocular lenses: overview of their capabilities, limitations, and clinical benefits. 24(3):215-217. [Editorial]
- Knorz MC. See Jendritza BB.
- Koivula A, Taube M, Zetterström C. Phakic refractive lens: two-year results. 24(5):507-515.
- Kojima T. See Yoshida Y.
- Konig K. See Wang BG.
- Kounis G. See Kymionis GD.
- Krueger RR, Mrochen M. Introduction to the Proceedings of the Third International Congress of Corneal Cross-Linking. 24(7):S713-S714. [Proceedings]
- Krueger RR, Ramos-Esteban JC, Kanellopoulos AJ. Staged intrastromal delivery of riboflavin with UVA cross-linking in advanced bullous keratopathy: laboratory investigation and first clinical case. 24(7):S730-S736. [Proceedings]
- Krueger RR, Rocha KM. Introduction to wavefront-optimized, wavefront-guided, and topography-guided customized ablation: fifth year in review. 24(4):S417-S418.
- Krueger RR, Trattler W, Yee R. Introduction to the Proceedings of the Sixth International Congress on Advanced Surface Ablation & SBK. 24(1):S55-S56. [Proceedings]
- Krueger RR. See Applegate RA.
- Krueger RR. See Iseli HP.
- Krueger RR. See Lin DTC.
- Krueger RR. See Ramos-Esteban JC.
- Krueger RR. See Rocha KM.
- Krueger RR. See Thornton I.
- Krueger RR. See Utz VM.
- Kuhlisch E. See Spoerl E.
- Kymionis GD, Diakonou VF, Charisis S, Pallikaris AI, Bouzoukis D, Yoo SH, Naoumidi I, Tsilimbaris MK. Effects of topical mitomycin C on the ciliary body and intraocular pressure after PRK: an experimental study. 24(6):633-638.
- Kymionis GD, Diakonou VF, Kounis G, Charisis S, Bouzoukis D, Ginis H, Yoo S, Tsilimbaris M, Pallikaris IG. Ocular rigidity evaluation after photorefractive keratectomy: an experimental study. 24(2):173-177. [Biomechanics]
- Kymionis GD. See Ide T.
- Lam DSC. See Cheng ACK.
- Lamoureux E. See Wittig-Silva C.
- Ltkany R. See Khalil M.

- Lau S. See Cheng ACK.
- Lazzarotto M. See Casprini F.
- Lee CS. See Lee JH.
- Lee DH. See Lee JH.
- Lee HK. See Chung SA.
- Lee HM. See Wang JC.
- Lee JH, Stulting RD, Lee DH, Lee CS, Kim WC, Kim EK Exacerbation of granular corneal dystrophy type II (Avellino corneal dystrophy) after LASEK. 24(1):39-45.
- Lepri F. See Casprini F.
- Lessell S. See Montezuma SR.
- Leung C. See Cheng ACK.
- Leung CKS. See Cheng ACK.
- Levinger S. See Barequet IS.
- Li J. See Liu Z.
- Li Y. See Liu Z.
- Lim KL. See Fam HB.
- Lim MCC, Chan TK. Central serous chorioretinopathy following LASIK for hyperopia. 24(6):651-652. [Report]
- Lima-Filho AAS. See Bottós KM.
- Lin DTC, Holland SP, Rocha KM, Krueger RR. Method for optimizing topography-guided ablation of highly aberrated eyes with the ALLEGRETTO WAVE excimer laser. 24(4):S439-S445.
- Lin JM, Chen WL, Chiang CC, Tsai YY. Ablation centration after active eye tracker-assisted LASIK and comparison of flying-spot and broad-beam laser. 24(4):371-376.
- Lin JT. Comparing anterior and posterior piggyback IOL power calculations in 2-optics and 3-optics systems. 24(7):665-666. [Letter]
- Lindsay RG. See Wittig-Silva C.
- Liu Y. See Warden L.
- Liu Z, Li Y, Cheng Z, Zhou F, Jiang H, Li J. Seven-year follow-up of LASIK for moderate to severe myopia. 24(9):935-940.
- Lohmann CP. See Wang BG.
- Lovatt ME. See Ursea R.
- Lubatschowski H. Overview of commercially available femtosecond lasers in refractive surgery. 24(1):S102-S107. [Proceedings]
- Lubatschowski H. See Schumacher S.
- MacRae S. See Yoon G.
- Maden A. See Yilmaz S.
- Malandrini A. See Casprini F.
- Maldonado MJ, Corcostegui I, García-Layana A, Salinas-Alaman A, Rutzen AR. Laser refractive surgery in a patient with a prepapillary arterial loop. 24(1):49-51. [Report]
- Malecaze F. See Couillet J.
- Maloney RK. See Jampaulo M.
- Marina-Verde C. See Gil-Cazorla R.
- Marotta G. See Cennamo G.
- Marshall J. See Durrie DS.
- Martin M. See Scheuerle AF.
- Martone G. See Casprini F.
- Mas D. See Illueca C.
- McCarey BE. See Dawson DG.
- McCarey BE. See Randleman JB.
- McCauley MB, Anderson DM, Johnson AJ. Posterior chamber Visian implantable collamer lens: stability and evaluation following traumatic grenade explosion. 24(6):648-651. [Report]
- McCulley JP. See Awwad ST.
- McCulley JP. See Petroll WM.
- McDonnell PJ. See Behrens A.
- Mearza AA, Muhtaseb M, Aslanides IM. Visual and refractive outcomes of LASIK with the SCHWIND ESIRIS and WaveLight ALLEGRETTO WAVE Eye-Q excimer lasers: a prospective, contralateral study. 24(9):885-890.
- Mertens EL, Sanders DR, Vitale PN. Custom-designed toric phakic intraocular lenses to correct high corneal astigmatism. 24(5):501-506.
- Mester U, Kaymak H. Comparison of the AcrySof IQ aspheric blue light filter and the AcrySof SA60AT intraocular lenses. 24(8):817-825.
- Mester U. See Kaymak H.
- Miller D. Anterior segment optical coherence tomography. 24(6):565. [Book Review]
- Miller D. Irregular astigmatism: diagnosis and treatment. 24(8):852. [Book Review]
- Miller D. Presbyopic lens surgery: a clinical guide to current technology. 24(7):756. [Book Review]
- Miraftaab M. See Hashemi H.
- Moghimi S. See Hashemi H.
- Mohammadi M. See Hashemi H.
- Mohan RR. See de Medeiros FW.
- Moilanen JAO. See Neira-Zalentein W.
- Montés-Micó R. See Alfonso JF.
- Montés-Micó R. See Cadarso L.
- Montés-Micó R. See Cerviño A.
- Montés-Micó R. See Charman WN.
- Montezuma SR, Lessell S, Pineda R. Optic neuropathy after ep LASIK. 24(2):204-208. [Report]
- Moore CR. See Kezirian GM.
- Mootha VV. See Petroll WM.
- Morton S. See Jendritza BB.
- Mosquera SA. See de Ortueta D.
- Mrochen M, Bueeler M, Donitzky C, Seiler T. Optical ray tracing for the calculation of optimized corneal ablation profiles in refractive treatment planning. 24(4):S446-S451.
- Mrochen M. See Krueger RR.
- Muftuoglu O. See Erdem U.
- Muhtaseb M. See Mearza AA.
- Muñoz G. See Javaloy J.
- Murat D. See Yamaguchi T.
- Mutani B. See de Sanctis U.
- Nader HB. See Bottós KM.
- Nakamura T. See Yoshida Y.
- Nalgirkar A. See Ursea R.
- Naoumidi I. See Kymionis GD.
- Narvaez J. See Waring GO III.
- Negishi K. See Yamaguchi T.
- Neira-Zalentein W, Moilanen JAO, Tuisku IS, Holopainen JM, Tervo TMT. Photorefractive keratectomy retreatment after LASIK. 24(7):710-712. [Report]
- Netto MV, Barreto J Jr, Santo R, Bechara S, Kara-Jose N, Wilson SE. Synergistic effect of ethanol and mitomycin C on corneal stroma. 24(6):626-632.
- Nitter TA. See Stojanovic A.
- Nosé W. See Fontes BM.
- O'Brien TP. See Ide T.
- O'Keefe M. See Kirwan C.

- Oberheide U. See Schumacher S.
- Ohtake Y. See Yamaguchi T.
- Ollero A. See Cadarso L.
- Ortiz D. See Illueca C.
- Ortiz D. See Pinelli R.
- Ott G. See Kaymak H.
- Ozkilic E. See Ertan A.
- Paik DC, Wen Q, Braunstein RE, Trokel SL. Short chain aliphatic  $\beta$ -nitro alcohols for corneoscleral cross-linking: corneal endothelial toxicity studies. 24(7):S741-S747. [Proceedings]
- Palacios A. See Alfonso JF.
- Pallikaris AI. See Kymionis GD.
- Pallikaris IG. See Kymionis GD.
- Palmer AM, Faiña PG, Albelda AE, Serrano MC, Saad DN, Céspedes MC. Visual function with bilateral implantation of monofocal and multifocal intraocular lenses: A prospective, randomized, controlled clinical trial. 24(3):257-264.
- Pantanelli S. See Yoon G.
- Papadia M. See Iester M.
- Park SH. See Jung SW.
- Park SH, Kim SY, Kim HI, Yang SW. Urrets-Zavalía syndrome following iris-claw phakic intraocular lens implantation. 24(9):959-961. [Report]
- Patel S, Alió JL, Feinbaum C. Comparison of Acri.Smart multifocal IOL, Crystalens AT-45 accommodative IOL, and Technovision presbyLASIK for correcting presbyopia. 24(3):294-299.
- Pellegrini G. See Vinciguerra P.
- Pereira T. See Vieira AC.
- Perente I. See Utine CA.
- Pérez J. See Illueca C.
- Pérez-Santonja JJ. See Javaloy J.
- Peters NT. OCT analysis of flap thickness. 24(2):117. [Letter]
- Petroll WM, Bowman RW, Cavanagh HD, Verity SM, Mootha VV, McCulley JP. Assessment of keratocyte activation following LASIK with flap creation using the IntraLase FS60 laser. 24(8):847-849. [Report]
- Peyman GA, Sanders DR, Battle JF, Féliz R, Cabrera G. Cyclosporine 0.05% ophthalmic preparation to aid recovery from loss of corneal sensitivity after LASIK. 24(4):337-343.
- Phillips S. See Waring GO III.
- Piers P. See Terwee T.
- Pillunat LE. See Spoerl E.
- Pineda R. See Montezuma SR.
- Pinelli R, Ortiz D, Simonetto A, Bacchi C, Sala E, Alió JL. Correction of presbyopia in hyperopia with a center-distance, paracentral-near technique using the Technolas 217z platform. 24(5):494-500.
- Pinelli R. More on peripheral presbyLASIK as a center-distance technique. 24(7):665. [Reply to Letter]
- Piñero DP. See Alió JL.
- Pita B. See Cadarso L.
- Pratzer KA. See Behrens A.
- Prescott C. See Blaylock JF.
- Purcell TL. See Ursea R.
- Pushpoth S. See Savant V.
- Qian Y. See Rocha KM.
- Rabinowitz YS. Metrics for keratoconus. 24(5):461. [Reply to Letter]
- Radhakrishnan H. See Charman WN.
- Raiskup-Wolf F. See Spoerl E.
- Rakpanichmanee T. See Srivannaboon S.
- Rama P. See Elsheikh A.
- Ramos-Esteban JC, Bamba S, Krueger RR. Tracking difficulties after femtosecond laser flap creation with the LADARVision excimer laser system. 24(9):953-956. [Report]
- Ramos-Esteban JC. See Krueger RR.
- Ramos-Esteban JC. See Rocha KM.
- Randleman JB, Dawson DG, Grossniklaus HE, McCarey BE, Edelhauser HF. Depth-dependent cohesive tensile strength in human donor corneas: implications for refractive surgery. 24(1):S85-S89. [Proceedings]
- Rao SK. See Cheng ACK.
- Rao SSK. See Cheng ACK.
- Regatieri CVS. See Bottós KM.
- Reilly C. See Caldwell M.
- Reinstein DZ, Archer TJ, Gobbe M, Silverman RH, Coleman DJ. Epithelial thickness in the normal cornea: three-dimensional display with Artemis very high-frequency digital ultrasound. 24(6):571-581.
- Reznik J, Salz JJ, Klimava A. Development of unilateral corneal ectasia after PRK with ipsilateral preoperative forme fruste keratoconus. 24(8):843-847. [Report]
- Riemann I. See Wang BG.
- Rivera RP. See Sanders DR.
- Rocha KM, Ramos-Esteban JC, Qian Y, Herekar S, Krueger RR. Comparative study of riboflavin-UVA cross-linking and "flash-linking" using surface wave elastometry. 24(7):S748-S751. [Proceedings]
- Rocha KM. See Krueger RR.
- Rocha KM. See Lin DTC.
- Rohrig B. See Hütz WW.
- Rosetta P. See Vinciguerra P.
- Rutzen AR. See Maldonado MJ.
- Ryu IKH. See Chung SA.
- Saad DN. See Palmer AM.
- Sala E. See Illueca C.
- Sala E. See Pinelli R.
- Salinas-Alaman A. See Maldonado MJ.
- Salz JJ. See Reznik J.
- Samimi D, Hamilton DR. Recalcitrant epithelial ingrowth in patients with compromised eyelid function. 24(5):544-546. [Report]
- Sánchez-Pina JM. See de Benito-Llopis L.
- Sanders DR. Anterior subcapsular opacities and cataracts 5 years after surgery in the Visian Implantable Collamer Lens FDA trial. 24(6):566-570.
- Sanders DR, Bernitsky DA, Harton PJ Jr, Rivera RP. The Visian myopic implantable collamer lens does not significantly affect axial length measurement with the IOLMaster. 24(9):957-959. [Report]
- Sanders DR, Sanders ML. Comparison of the toric implantable collamer lens and custom ablation LASIK for myopic astigmatism. 24(8):773-778.
- Sanders DR. See Mertens EL.
- Sanders DR. See Peyman GA.
- Sanders ML. See Sanders DR.
- Santo R. See Netto MV.
- Savant V, Chavan R, Pushpoth S, Ilango B. Comparability and intra-/interobserver reliability of anterior chamber depth measurements with the Pentacam and IOLMaster. 24(6):615-618.

- Savini G, Barboni P, Zanini M. Calculating the keratometric refractive index. 24(1):10-11. [Reply to Letter]
- Savini G, Hoffer KJ. Corneal power after myopic LASIK. 24(8):769. [Letter]
- Schanzer C. See Waring GO III.
- Schanzlin DJ. See Ursea R.
- Scheuerle AF, Martin M, Voelcker HE, Auffarth G. Undetected development of glaucoma after radial keratotomy. 24(1):51-54. [Report]
- Schmickler S. Postoperative changes in refraction and anterior chamber depth in different multifocal intraocular lenses. 24(3):306-308. [Report]
- Schor CM, Bharadwaj SR. Adaptive calibration of dynamic accommodation—implications for accommodating intraocular lenses. 24(9):984-990. [Proceedings]
- Schor P. See Bottós KM.
- Schubert H. See Wang BG.
- Schumacher S, Fromm M, Oberheide U, Gerten G, Wegener A, Lubatschowski H. In vivo application and imaging of intralenticular femtosecond laser pulses for the restoration of accommodation. 24(9):991-995. [Proceedings]
- Schwendeman F. See Eser I.
- Schwendeman FJ. See Stahl JE.
- Schwiegerling J, Choi J. Application of the polychromatic defocus transfer function to multifocal lenses. 24(9):965-969. [Proceedings]
- Schwiegerling J. See Choi J.
- Seiler T. See Iseli HP.
- Seiler T. See Mrochen M.
- Serrano MC. See Palmer AM.
- Shimizu K. See Iida Y.
- Shimizu K. See Kamiya K.
- Si Z. See Blaylock JF.
- Silverman RH. See Reinstejn DZ.
- Simonetto A. See Pinelli R.
- Sinhá S. See de Medeiros FW.
- Slade SG. See Durrie DS.
- Snibson GR. See Wittig-Silva C.
- Sobaci G. See Erdem U.
- Souza DC. See Ghanem VC.
- Souza GC. See Ghanem VC.
- Speaker MG. See Khalil M.
- Spoerl E, Raiskup-Wolf F, Kuhlisch E, Pillunat LE. Cigarette smoking is negatively associated with keratoconus. 24(7):S737-S740. [Proceedings]
- Spoerl E. See Iseli HP.
- Srinivasan S, Drake A, Herzig S. Photorefractive keratectomy with 0.02% mitomycin C for treatment of residual refractive errors after LASIK. 24(1):S64-S67. [Proceedings]
- Srivannaboon S, Rakpanichmanee T, Cheng ACK, Fam HB. Estimation of posterior corneal power for IOL calculation after myopic LASIK. 24(9):946-951. [Biomechanics]
- Stahl JE, Durrie DS, Schwendeman FJ, Boghossian AJ. OCT analysis of flap thickness. 24(2):117-119. [Reply to Letter]
- Stahl JE, Durrie DS, Schwendeman FJ, Boghossian AJ. Planar flaps with the Carriazo-Pendular microkeratome. 24(4):322-323. [Reply to Letter]
- Stahl JE. See Eser I.
- Stark WJ. See Behrens A.
- Stevens G. See Waring GO III.
- Stojanovic A, Wang L, Jankov M, Nitter TA, Wang Q. Wavefront optimized versus custom-Q treatments in surface ablation for myopic astigmatism with the WaveLight ALLEGRETTO laser. 24(8):779-789.
- Stonecipher KG, Kezirian GM. Wavefront-optimized versus wavefront-guided LASIK for myopic astigmatism with the ALLEGRETTO WAVE: three-month results of a prospective FDA trial. 24(4):S424-S430.
- Stonecipher KG. See Kezirian GM.
- Stulting RD. See Lee JH.
- Sullivan LJ. See Wittig-Silva C.
- SurgiVision® Consultants Inc WaveLight Investigator Group. See Kezirian GM.
- Suto C. See de Medeiros FW.
- Sutton G, Hodge C. Accuracy and precision of LASIK flap thickness using the IntraLase femtosecond laser in 1000 consecutive cases. 24(8):802-806.
- Suzuki M. See Iida Y.
- Sverdrup L. See Warden L.
- Tan BU. See Ursea R.
- Tanchel N. See Waring GO III.
- Taube M. See Koivula A.
- Tchah H. See Choi CY.
- Tervo T. See Ambrósio R Jr.
- Tervo TM. See Tuisku IS.
- Tervo TMT. See Neira-Zalentein W.
- Terwee T, Weeber H, van der Mooren M, Piers P. Visualization of the retinal image in an eye model with spherical and aspheric, diffractive, and refractive multifocal intraocular lenses. 24(3):223-232.
- Teus MA. See de Benito-Llopis L.
- Teus MA. See Gil-Cazorla R.
- Thibos LN. Where is the optimum far-point for a presbyopic eye? 24(9):970-975. [Proceedings]
- Thornton I, Xu M, Krueger RR. Comparison of standard (0.02%) and low dose (0.002%) mitomycin C in the prevention of corneal haze following surface ablation for myopia. 24(1):S68-S76. [Proceedings]
- Tomas JD. See Uceda-Montanes A.
- Tosi GM. See Casprini F.
- Trattler W. See Krueger RR.
- Traverso CE. See Iester M.
- Trokel SL. See Paik DC.
- Tsai YY. See Lin JM.
- Tsilimbaris M. See Kymionis GD.
- Tsubota K. See Yamaguchi T.
- Tuisku IS, Tervo TM, Belmonte C. Tricyclic antidepressants: potential therapeutic alternatives for treatment of dry eye symptoms after LASIK. 24(8):771-772. [Reply to Letter]
- Tuisku IS. See Neira-Zalentein W.
- Uceda-Montanes A, Tomás JD, Alió JL. Correction of severe ectasia after LASIK with intracorneal ring segments. 24(4):408-411. [Report]
- Ursea R, Purcell TL, Tan BU, Nalgirkar A, Lovaton ME, Ehrenhaus MP, Schanzlin DJ. The effect of cyclosporine A (Restasis) on recovery of visual acuity following LASIK. 24(5):473-476.
- Utine CA, Cakir H, Egemenoglu A, Perente I. LASIK in children with hyperopic anisometropic amblyopia. 24(5):464-472.
- Utz VM, Krueger RR. Management of irregular astigmatism following rotationally disoriented free cap after LASIK. 24(4):383-391.

- van der Mooren M. See Terwee T.
- Velarde GC. See Fontes BM.
- Venter J. Wavefront-guided custom ablation for myopia using the NIDEK NAVEX laser system. 24(5):487-493.
- Venturino G. See Iester M.
- Verity SM. See Petroll WM.
- Vidal MT. See Javaloy J.
- Vieira AC, Pereira T, de Freitas D. Late-onset infections after LASIK. 24(4):411-413. [Report]
- Viese JMZ. See Ghanem VC.
- Vinciguerra P, Albè E, Rosetta P, Di Iorio E, Pellegrini G. Custom phototherapeutic keratectomy and autologous fibrin-cultured limbal stem cell autografting: a combined approach. 24(4):323-324. [Letter]
- Vitale PN. See Mertens EL.
- Voelcker HE. See Scheuerle AF.
- Vossmerbaeumer U, Jonas JB. Regularity of human corneal flaps prepared by femtosecond laser technology. 24(6):645-648. [Report]
- Vossmerbaeumer U. See Knorz MC.
- Wallau AD, Campos M. Photorefractive keratectomy with mitomycin C versus LASIK in custom surgeries for myopia: a bilateral prospective randomized clinical trial. 24(4):326-336.
- Wang BG, Lohmann CP, Riemann I, Schubert H, Halbhuber KJ, König K. Multiphoton-mediated corneal flap generation using the 80 MHz nanjoule femtosecond near-infrared laser. 24(8):833-839.
- Wang F. See Yu J.
- Wang JC, Bunce C, Lee HM. Intraoperative corneal thickness measurement using optical coherence pachymetry and corneo-gage plus ultrasound pachymetry. 24(6):610-614.
- Wang L. See Stojanovic A.
- Wang Q. See Stojanovic A.
- Warden L, Liu Y, Binder PS, Dreher AW, Sverdrup L. Performance of a new binocular wavefront aberrometer based on a self-imaging diffractive sensor. 24(2):188-196. [New Technology]
- Waring GO III, Fant B, Stevens G, Phillips S, Fischer J, Tanchel N, Schanzer C, Narvaez J, Chayet A. Laser in situ keratomileusis for spherical hyperopia and hyperopic astigmatism using the NIDEK EC-5000 excimer laser. 24(2):123-136.
- Waring GO III. Excellence. 24(7):662-663. [Editorial]
- Waring GO III. Neo-neologisms. 24(1):7-8. [Editorial]
- Waring GO III. Troutman award for editorial excellence—16 years of inspired contributions. 24(9):863-864. [Editorial]
- Waring GO IV, Durrie DS. Emerging trends for procedure selection in contemporary refractive surgery: consecutive review of 200 cases from a single center. 24(4):S419-S423.
- Warmerdam D. See Awwad ST.
- Weber SLP. See Ghanem VC.
- Weeber H. See Terwee T.
- Wegener A. See Schumacher S.
- Wen Q. See Paik DC.
- Whiting M. See Wittig-Silva C.
- Wiedemann P. See Iseli HP.
- Wilson SE. See Ambrósio R Jr.
- Wilson SE. See de Medeiros FW.
- Wilson SE. See Netto MV.
- Wittig-Silva C, Whiting M, Lamoureux E, Lindsay RG, Sullivan LJ, Snibson GR. A randomized controlled trial of corneal collagen cross-linking in progressive keratoconus: preliminary results. 24(7):S720-S725. [Proceedings]
- Wong A. See Cheng ACK.
- Wong AL. See Cheng ACK.
- Wyler D. See Camellin M.
- Xu M. See Thornton I.
- Yamaguchi T, Murat D, Kimura I, Negishi K, Yuki K, Tsubota K, Ohtake Y. Diagnosis of steroid-induced glaucoma after photorefractive keratectomy. 24(4):413-415. [Report]
- Yang SW. See Park SH.
- Yee R. See Krueger RR.
- Yildiz TF. See Akin T.
- Yilmaz S, Yuksel T, Maden A. Corneal topographic changes after four types of pterygium surgery. 24(2):160-165.
- Yoo S. See Kymionis GD.
- Yoo SH. See Ide T.
- Yoon G, Pantanelli S, MacRae S. Comparison of Zernike and Fourier wavefront reconstruction algorithms in representing corneal aberration of normal and abnormal eyes. 24(6):582-590.
- Yoshida N. See Yoshida Y.
- Yoshida Y, Nakamura T, Hara S, Yoshida N, Kojima T, Ichikawa K. Topography-guided custom ablation for irregular corneal astigmatism using the NIDEK NAVEX laser system. 24(1):24-32.
- Yu G. See Khalil M.
- Yu J, Chen H, Wang F. Patient satisfaction and visual symptoms after wavefront-guided and wavefront-optimized LASIK with the WaveLight platform. 24(5):477-486.
- Yuki K. See Yamaguchi T.
- Yuksel T. See Yilmaz S.
- Zakrzewski PA. See Izquierdo L Jr.
- Zanini M. See Savini G.
- Zetterström C. See Koivula A.
- Zhou F. See Liu Z.
- Zhou XT. See Cheng ZY.

#### TITLE INDEX

- A comparative analysis of intraocular lens power calculation methods after myopic excimer laser surgery. Fam HB, Lim KL. 24(4):355-360.
- A prospective bilateral comparison of epi-LASIK and LASEK for myopia. Hondur A, Bilgihan K, Hasanreisoglu B. 24(9):928-934.
- A randomized controlled trial of corneal collagen cross-linking in progressive keratoconus: preliminary results. Wittig-Silva C, Whiting M, Lamoureux E, Lindsay RG, Sullivan LJ, Snibson GR. 24(7):S720-S725. [Proceedings]
- A retrospective comparison of LASIK outcomes for myopia and myopic astigmatism with conventional NIDEK versus wavefront-guided VISX and Alcon platforms. Dougherty PJ, Bains HS. 24(9):891-896.
- A systematic review of Ferrara's ring in the treatment of keratoconus. El Dib RP, de Freitas D. 24(9):865-866. [Letter]
- Ablation centration after active eye tracker-assisted LASIK and comparison of flying-spot and broad-beam laser. Lin JM, Chen WL, Chiang CC, Tsai YY. 24(4):371-376.
- Accuracy and precision of LASIK flap thickness using the IntraLase femtosecond laser in 1000 consecutive cases. Sutton G, Hodge C. 24(8):802-806.
- Adaptive calibration of dynamic accommodation—implications for accommodating intraocular lenses. Schor CM, Bharadwaj SR. 24(9):984-990. [Proceedings]
- Anterior segment optical coherence tomography. Miller DM. 24(6):565. [Book Review]

- Anterior subcapsular opacities and cataracts 5 years after surgery in the Visian Implantable Collamer Lens FDA trial. Sanders DR. 24(6):566-570.
- Application of the polychromatic defocus transfer function to multifocal lenses. Schwiegerling J, Choi J. 24(9):965-969. [Proceedings]
- Assessment of keratocyte activation following LASIK with flap creation using the IntraLase FS60 laser. Petroll WM, Bowman RW, Cavanagh HD, Verity SM, Mootha VV, McCulley JP. 24(8):847-849. [Report]
- Association between ocular dominance and refraction. Eser I, Durrie DS, Schwendeman F, Stahl JE. 24(7):685-689.
- Biomechanical and wound healing characteristics of corneas after excimer laser keratorefractive surgery: is there a difference between advanced surface ablation and sub-Bowman's keratomileusis? Dawson DG, Grossniklaus HE, McCarey BE, Edelhauser HF. 24(1):S90-S96. [Proceedings]
- Calculating the keratometric refractive index. Borrelli M, Irregolare C. 24(1):10. [Letter]
- Calculating the keratometric refractive index. Savini G, Barboni P, Zanini M. 24(1):10-11. [Reply to Letter]
- Central corneal thickness measurements by ultrasound, Orbscan II, and Visante OCT after LASIK for myopia. Cheng ACK, Rao SK, Lau S, Leung CKS, Lam DSC. 24(4):361-365.
- Central serous chorioretinopathy following LASIK for hyperopia. Lim MCC, Chan TK. 24(6):651-652. [Report]
- Cigarette smoking is negatively associated with keratoconus. Spoerl E, Raiskup-Wolf F, Kuhlisch E, Pillunat LE. 24(7):S737-S740. [Proceedings]
- Comparability and intra-/interobserver reliability of anterior chamber depth measurements with the Pentacam and IOLMaster. Savant V, Chavan R, Pushpoth S, Ilango B. 24(6):615-618.
- Comparative study of riboflavin-UVA cross-linking and "flash-linking" using surface wave elastometry. Rocha KM, Ramos-Esteban JC, Qian Y, Herekar S, Krueger RR. 24(7):S748-S751. [Proceedings]
- Comparing anterior and posterior piggyback IOL power calculations in 2-optics and 3-optics systems. Lin JT. 24(7):665-666. [Letter]
- Comparison between LASEK with mitomycin C and LASIK for the correction of myopia of  $-7.00$  to  $-13.75$  D. de Benito-Llopis L, Teus MA, Sánchez-Pina JM. 24(5):516-523.
- Comparison of Acri.Smart multifocal IOL, Crystalens AT-45 accommodative IOL, and Technovision presbyLASIK for correcting presbyopia. Patel S, Alió JL, Feinbaum C. 24(3):294-299.
- Comparison of corneal deposits after LASIK and PRK in eyes with granular corneal dystrophy type II. Kim T, Kim T, Kim SW, Kim EK. 24(4):392-395.
- Comparison of flap adhesion strength using the Amadeus microkeratome and the IntraLase iFS femtosecond laser in rabbits. Knorz MC, Vossmerbaeumer U. 24(9):875-878.
- Comparison of higher order aberrations measured by NIDEK OPD-Scan dynamic skiascopy and Zeiss WASCAs Hartmann-Shack aberrometers. Cerviño A, Hosking SL, Montés-Micó R. 24(8):790-796.
- Comparison of silicone and non-silicone hydrogel soft contact lenses used as a bandage after LASEK. Gil-Cazorla R, Teus MA, Arranz-Márquez E. 24(2):199-203. [Report]
- Comparison of standard (0.02%) and low dose (0.002%) mitomycin C in the prevention of corneal haze following surface ablation for myopia. Thornton I, Xu M, Krueger RR. 24(1):S68-S76. [Proceedings]
- Comparison of techniques for corneal power assessment after myopic LASIK without the use of preoperative data. Cheng ACK, Rao SSK, Lau S, Wong A, Lam DSC. 24(5):539-543.
- Comparison of the AcrySof IQ aspheric blue light filter and the AcrySof SA60AT intraocular lenses. Mester U, Kaymak H. 24(8):817-825.
- Comparison of the toric implantable collamer lens and custom ablation LASIK for myopic astigmatism. Sanders DR, Sanders ML. 24(8):773-778.
- Comparison of Zernike and Fourier wavefront reconstruction algorithms in representing corneal aberration of normal and abnormal eyes. Yoon G, Pantanelli S, MacRae S. 24(6):582-590.
- Complications of sub-Bowman's keratomileusis with a femtosecond laser in 3009 eyes. Chang JSM. 24(1):S97-S101. [Proceedings]
- Contrast sensitivity and higher order aberrations in eyes implanted with AcrySof IQ SN60WF and AcrySof SN60AT intraocular lenses. Awwad ST, Warmerdam D, Bowman RW, Dwarakanathan S, Cavanagh HD, McCulley JP. 24(6):619-625.
- Corneal biomechanical metrics in eyes with refraction of  $-19.00$  to  $+9.00$  D in healthy Brazilian patients. Fontes BM, Ambrósio R Jr, Alonso RS, Jardim D, Velarde GC, Nosé W. 24(9):941-945. [Biomechanics]
- Corneal power after myopic LASIK. Savini G, Hoffer KJ. 24(8):769. [Letter]
- Corneal power after myopic LASIK. Cheng ACK, Rao SSK, Lau S, Wong A, Lam DSC. 24(8):770. [Reply to Letter]
- Corneal topographic changes after four types of pterygium surgery. Yilmaz S, Yuksel T, Maden A. 24(2):160-165.
- Correction of presbyopia in hyperopia with a center-distance, paracentral-near technique using the Technolas 217z platform. Pinelli R, Ortíz D, Simonetto A, Bacchi C, Sala E, Alió JL. 24(5):494-500.
- Correction of severe ectasia after LASIK with intracorneal ring segments. Uceda-Montanes A, Tomás JD, Alió JL. 24(4):408-411. [Report]
- Custom phototherapeutic keratectomy and autologous fibrin-cultured limbal stem cell autografting: a combined approach. Vinciguerra P, Albè E, Rosetta P, Di Iorio E, Pellegrini G. 24(4):323-324. [Letter]
- Custom-designed toric phakic intraocular lenses to correct high corneal astigmatism. Mertens EL, Sanders DR, Vitale PN. 24(5):501-506.
- Customized topography-guided photorefractive keratectomy with the MEL-70 platform and mitomycin C to correct hyperopia after radial keratotomy. Ghanem RC, Ghanem VC, de Souza DC, Karajósé N, Ghanem EA. 24(9):911-922.
- Cyclosporine 0.05% ophthalmic preparation to aid recovery from loss of corneal sensitivity after LASIK. Peyman GA, Sanders DR, Batlle JF, Feliz R, Cabrera G. 24(4):337-343.
- Depth-dependent cohesive tensile strength in human donor corneas: implications for refractive surgery. Randleman JB, Dawson DG, Grossniklaus HE, McCarey BE, Edelhauser HF. 24(1):S85-S89. [Proceedings]
- Detection of an abnormally thick LASIK flap with anterior segment OCT imaging prior to planned LASIK retreatment surgery. Izquierdo L Jr, Henriquez MA, Zakrzewski PA. 24(2):197-199. [Report]
- Development of unilateral corneal ectasia after PRK with ipsilateral preoperative forme fruste keratoconus. Reznik J, Salz JJ, Klimava A. 24(8):843-847. [Report]
- Diagnosis of steroid-induced glaucoma after photorefractive keratectomy. Yamaguchi T, Murat D, Kimura I, Negishi K, Yuki K, Tsubota K, Ohtake Y. 24(4):413-415. [Report]
- Dynamics of small-incision clear cornea wounds after phacoemulsification surgery using optical coherence tomography in the early postoperative period. Behrens A, Stark WJ, Pratzler KA, McDonnell PJ. 24(1):46-49. [Report]

- Effect of age on outcomes in patients with keratoconus treated by Intacs using a femtosecond laser. Ertan A, Ozkiloglu E. 24(7):690-695.
- Effect of flap thickness on higher order wavefront aberrations induced by LASIK: a bilateral study. Cheng ZY, He JC, Zhou XT, Chu RY. 24(5):524-529.
- Effect of thin femtosecond LASIK flaps on corneal sensitivity and tear function. Barequet IS, Hirsh A, Levinger S. 24(9):897-902.
- Effectiveness of cultured human keratinocyte onlays on epithelial healing and clinical outcome after photorefractive keratectomy. Chung SA, Kim EK, Ryu IKH, Kim JK, Lee HK. 24(8):826-832.
- Effects of topical mitomycin C on the ciliary body and intraocular pressure after PRK: an experimental study. Kymionis GD, Diakonis VF, Charisis S, Pallikaris AI, Bouzoukis DI, Yoo SH, Naoumidis I, Tsilimbaris MK. 24(6):633-638.
- Effects of topical nepafenac on corneal epithelial healing time and postoperative pain after PRK: a bilateral, prospective, randomized, masked trial. Caldwell M, Reilly C. 24(4):377-382.
- Efficacy and safety of blue-light scleral cross-linking. Iseli HP, Spoerl E, Wiedemann P, Krueger RR, Seiler T. 24(7):S752-S755. [Proceedings]
- Emerging trends for procedure selection in contemporary refractive surgery: consecutive review of 200 cases from a single center. Waring GO IV, Durrie DS. 24(4):S419-S423.
- Epi-LASIK versus epi-LASEK. Camellin M, Wyler D. 24(1):S57-S63. [Proceedings]
- Epithelial thickness in the normal cornea: three-dimensional display with Artemis very high-frequency digital ultrasound. Reinstein DZ, Archer TJ, Gobbe M, Silverman RH, Coleman DJ. 24(6):571-581.
- Estimation of posterior corneal power for IOL calculation after myopic LASIK. Srivannaboon S, Rakpanichmanee T, Cheng ACK, Fam HB. 24(9):946-951. [Biomechanics]
- Exacerbation of granular corneal dystrophy type II (Avellino corneal dystrophy) after LASEK. Lee JH, Stulting RD, Lee DH, Lee CS, Kim WC, Kim EK. 24(1):39-45.
- Excellence. Waring GO III. 24(7):662-663. [Editorial]
- Experimental assessment of corneal anisotropy. Elsheikh A, Brown M, Alhasso D, Rama P, Campanelli M, Garway-Heath D. 24(2):178-187. [Biomechanics]
- Factors influencing flap and INTACS decentration after femtosecond laser application in normal and keratoconic eyes. Ertan A, Karacal H. 24(8):797-801.
- Four-year postoperative results of the US ALLEGRETTO WAVE clinical trial for the treatment of hyperopia. Kezirian GM, Moore CR, Stonecipher KG, SurgiVision® Consultants Inc WaveLight Investigator Group. 24(4):S431-S438.
- Haze development after photorefractive keratectomy: mechanical vs ethanol epithelial removal in rabbits. de Medeiros FW, Mohan RR, Suto C, Sinhá S, Bonilha VL, Chaurasia SS, Wilson SE. 24(9):923-927.
- Immunofluorescence confocal microscopy of porcine corneas following collagen cross-linking treatment with riboflavin and ultraviolet A. Bottós KM, Dreyfuss JL, Regatieri CVS, Lima-Filho AAS, Schor P, Nader HB, Chamon W. 24(7):S715-S719. [Proceedings]
- In vivo application and imaging of intralenticular femtosecond laser pulses for the restoration of accommodation. Schumacher S, Fromm M, Oberheide U, Gerten G, Wegener A, Lubatschowski H. 24(9):991-995. [Proceedings]
- Inadequate results for microkeratome-assisted additive stromal keratoplasty for management of keratoconus. Couillet J, Fournie P, Malecaze F, Arné JL. 24(2):166-172.
- Influence of age on ocular wavefront aberration changes with accommodation. Iida Y, Shimizu K, Ito M, Suzuki M. 24(7):696-701.
- Intacs implantation with sequential collagen cross-linking treatment in postoperative LASIK ectasia. Kamburoglu G, Ertan A. 24(7):S726-S729. [Proceedings]
- Intermediate vision and reading speed with Array, Tecnis, and ReSTOR intraocular lenses. Hütz WW, Eckhardt HB, Röhrig B, Grolmus R. 24(3):251-256.
- Intraocular comparison of the effect of training on visual performance with ReSTOR and Tecnis diffractive multifocal IOLs. Kaymak H, Fahle M, Ott G, Mester U. 24(3):287-293.
- Intraocular lens tilt and decentration after phacoemulsification surgery in patients with and without primary open angle glaucoma. Akin T, Aykan U, Yildiz TF, Karadayi K, Bilge AH. 24(9):865. [Letter]
- Intraoperative corneal thickness measurement using optical coherence pachymetry and corneo-gage plus ultrasound pachymetry. Wang JC, Bunce C, Lee HM. 24(6):610-614.
- Intrasubject corneal thickness asymmetry. Khachikian SS, Belin MW, Ciolino JB. 24(6):606-609.
- Introduction to the Proceedings of the 9th International Congress of Wavefront and Presbyopic Refractive Corrections. Applegate RA, Krueger RR. 24(9):963-964. [Proceedings]
- Introduction to the Proceedings of the Sixth International Congress on Advanced Surface Ablation & SBK. Krueger RR, Trattler W, Yee R. 24(1):S55-S56. [Proceedings]
- Introduction to the Proceedings of the Third International Congress of Corneal Cross-Linking. Krueger RR, Mrochen M. 24(7):S713-S714. [Proceedings]
- Introduction to wavefront-optimized, wavefront-guided, and topography-guided customized ablation: fifth year in review. Krueger RR, Rocha KM. 24(4):S417-S418.
- Irregular astigmatism: diagnosis and treatment. Miller D. 24(8):852. [Book Review]
- LASIK in children with hyperopic anisometropic amblyopia. Utine CA, Cakir H, Egemenoglu A, Perente I. 24(5):464-472.
- LASIK-associated dry eye and neurotrophic epitheliopathy: pathophysiology and strategies for prevention and treatment. Ambrósio R Jr, Tervo T, Wilson SE. 24(4):396-407. [Review]
- Lack of progression of ectasia seven years after LASIK in a highly myopic keratoconic eye. Jampaulo M, Maloney RK. 24(7):707-709. [Report]
- Laser in situ keratomileusis for spherical hyperopia and hyperopic astigmatism using the NIDEK EC-5000 excimer laser. Waring GO III, Fant B, Stevens G, Phillips S, Fischer J, Tanchel N, Schanzer C, Narvaez J, Chayet A. 24(2):123-136.
- Laser refractive surgery in a patient with a prepapillary arterial loop. Maldonado MJ, Corcóstegui I, García-Layana A, Salinas-Alaman A, Rutzen AR. 24(1):49-51. [Report]
- Late-onset infections after LASIK. Vieira AC, Pereira T, de Freitas D. 24(4):411-413. [Report]
- Long-term endothelial cell loss after traumatic dislocation and repositioning of artisan phakic IOL. de Sanctis U, Mutani B, Grignolo FM. 24(5):546-548.
- Long-term experience with mixing and matching refractive Array and diffractive CeeOn multifocal intraocular lenses. Gunenc U, Celik L. 24(3):233-242.
- Management of irregular astigmatism following rotationally disoriented free cap after LASIK. Utz VM, Krueger RR. 24(4):383-391.
- Mathematical properties of asphericity: a method to calculate with asphericities. Calossi A. 24(2):121. [Reply to Letter]
- Mathematical properties of asphericity: a method to calculate with asphericities. de Ortueta D, Mosquera SA. 24(2):119-121. [Letter]
- Measure of keratoconus progression using Orbscan II. Kim H, Joo CK. 24(6):600-605.

- Measurement of intraocular pressure in LASIK and LASEK patients using the Reichert Ocular Response Analyzer and Goldmann applanation tonometry. Kirwan C, O'Keefe M. 24(4):366-370.
- Measurement of LASIK flap thickness with anterior segment optical coherence tomography. Cheng ACK, Ho T, Lau S, Wong AL, Leung C, Lam DSC. 24(9):879-884.
- Mechanical corneal epithelium scraping and ethanol treatment up-regulate cytokine gene expression differently in rabbit cornea. Chang SW, Chou SF, Chuang JL. 24(2):150-159.
- Method for optimizing topography-guided ablation of highly aberrated eyes with the ALLEGRETTO WAVE excimer laser. Lin DTC, Holland SP, Rocha KM, Krueger RR. 24(4):S439-S445.
- Metrics for keratoconus. Boxer Wachler B. 24(5):460. [Letter]
- Metrics for keratoconus. Rabinowitz YS. 24(5):461. [Reply to Letter]
- Microbial keratitis after INTACS implantation with loose suture. Hashemi H, Ghaffari R, Mohammadi M, Moghimi S, Miraftaab M. 24(5):551-552.
- More on peripheral presbyLASIK as a center-distance technique. Pinelli R. 24(7):665. [Reply to Letter]
- Multifocal corneal ablation for hyperopic presbyopes. Jung SW, Kim MJ, Park SH, Joo CK. 24(9):903-910.
- Multifocal intraocular lenses: overview of their capabilities, limitations, and clinical benefits. Knorz MC. 24(3):215-217. [Editorial]
- Multiphoton-mediated corneal flap generation using the 80 MHz nanosecond femtosecond near-infrared laser. Wang BG, Lohmann CP, Riemann I, Schubert H, Halbhauer KJ, König K. 24(8):833-839.
- Myopic phakic STAAR collamer posterior chamber intraocular lenses for keratoconus. Alfonso JF, Palacios A, Montés-Micó R. 24(9):867-874.
- Neo-neologisms. Waring GO III. 24(1):7-8. [Editorial]
- OCT analysis of flap thickness. Peters NT. 24(2):117. [Letter]
- OCT analysis of flap thickness. Stahl JE, Durrie DS, Schwendeman FJ, Boghossian AJ. 24(2):117-119. [Reply to Letter]
- Ocular rigidity evaluation after photorefractive keratectomy: an experimental study. Kymionis GD, Diakonis VF, Kounis G, Charisis S, Bouzoukis D, Ginis H, Yoo S, Tsilimbaris M, Pallikaris IG. 24(2):173-177. [Biomechanics]
- Optic neuropathy after epi-LASIK. Montezuma SR, Lessell S, Pineda R. 24(2):204-208. [Report]
- Optical aberrations in pseudophakic eyes after 2.5-mm Nd:YAG laser capsulotomy for posterior capsule opacification. Casprini F, Balestrazzi A, Tosi GM, Lazzarotto M, Malandrini A, Lepri F, Martone G, Caporossi T, Caporossi A. 24(7):702-706.
- Optical performance measurement and night driving simulation of ReSTOR, ReZoom, and Tecnis multifocal intraocular lenses in a model eye. Choi J, Schwiegerling J. 24(3):218-222.
- Optical ray tracing for the calculation of optimized corneal ablation profiles in refractive treatment planning. Mrochen M, Bueeler M, Donitzky C, Seiler T. 24(4):S446-S451.
- Overview of commercially available femtosecond lasers in refractive surgery. Lubatschowski H. 24(1):S102-S107. [Proceedings]
- Patient satisfaction and visual symptoms after wavefront-guided and wavefront-optimized LASIK with the WaveLight platform. Yu J, Chen H, Wang F. 24(5):477-486.
- Performance of a new binocular wavefront aberrometer based on a self-imaging diffractive sensor. Warden L, Liu Y, Binder PS, Dreher AW, Sverdrup L. 24(2):188-196. [New Technology]
- Phakic refractive lens (Medennium) for correction of +4.00 to +6.00 diopters: 1-year follow-up. Gil-Cazorla R, Teus MA, Arranz-Marquez E, Marina-Verde C. 24(4):350-354.
- Phakic refractive lens: two-year results. Koivula A, Taube M, Zetterström C. 24(5):507-515.
- Phakic toric implantable collamer lens implantation for the correction of high myopic astigmatism in eyes with keratoconus. Kamiya K, Shimizu K, Ando W, Asato Y, Fujisawa T. 24(8):840-842. [Report]
- Photorefractive keratectomy and butterfly laser epithelial keratomileusis: a prospective, contralateral study. Ghanem VC, Kara-José N, Ghanem RC, Coral SA. 24(7):671-684.
- Photorefractive keratectomy retreatment after LASIK. Neira-Zalentein W, Moilanen JAO, Tuisku IS, Holopainen JM, Tervo TMT. 24(7):710-712. [Report]
- Photorefractive keratectomy with 0.02% mitomycin C for treatment of residual refractive errors after LASIK. Srinivasan S, Drake A, Herzig S. 24(1):S64-S67. [Proceedings]
- Photorefractive keratectomy with mitomycin C versus LASIK in custom surgeries for myopia: a bilateral prospective randomized clinical trial. Wallau AD, Campos M. 24(4):326-336.
- Planar flaps with the Carriazo-Pendular microkeratome. de Ortueta D. 24(4):322. [Letter]
- Planar flaps with the Carriazo-Pendular microkeratome. Stahl JE, Durrie DS, Schwendeman FJ, Boghossian AJ. 24(4):322-323. [Reply to Letter]
- Posterior chamber Visian implantable collamer lens: stability and evaluation following traumatic grenade explosion. McCauley MB, Anderson DM, Johnson AJ. 24(6):648-651. [Report]
- Posterior corneal curvature measurements with peripheral fitting zones before and after myopic LASIK using Orbscan II. Cheng ACK, Ho T, Rao SK, Lau S, Lam DSC. 24(8):807-810.
- Postoperative changes in refraction and anterior chamber depth in different multifocal intraocular lenses. Schmickler S. 24(3):306-308. [Report]
- Postoperative optical aberrations in eyes implanted with AcrySof spherical and aspheric intraocular lenses. Cadarso L, Iglesias A, Ollero A, Pita B, Montés-Micó R. 24(8):811-816.
- Presbyopic lens surgery: a clinical guide to current technology. Miller D. 24(7):756. [Book Review]
- PRK and butterfly LASEK: prospective, randomized, contralateral eye comparison of epithelial healing and ocular discomfort. Ghanem VC, Souza GC, Souza DC, Viese JMZ, Weber SLP, Kara-José N. 24(6):591-599.
- Problems in the measurement of wavefront aberration for eyes implanted with diffractive bifocal and multifocal intraocular lenses. Charman WN, Montés-Micó R, Radhakrishnan H. 24(3):280-286.
- Prospective evaluation of intraocular lens calculation after myopic refractive surgery. Khalil M, Chokshi A, Latkany R, Speaker MG, Yu G. 24(1):33-38.
- Pseudoaccommodation and visual acuity with Technovision presbyLASIK and a theoretical simulated Array® multifocal intraocular lens. Illueca C, Alió JL, Mas D, Ortiz D, Pérez J, Espinosa J, Sala E. 24(4):344-349.
- Pupil center shift relative to the coaxially sighted corneal light reflex under natural and pharmacologically dilated conditions. Erdem U, Muftuoglu O, Gundogan FC, Sobaci G, Bayer A. 24(5):530-538.
- Recalcitrant epithelial ingrowth in patients with compromised eyelid function. Samimi D, Hamilton DR. 24(5):544-546. [Report]
- Refractive lens exchange with the diffractive multifocal Tecnis ZM900 intraocular lens. Goes FJ. 24(3):243-250.
- Regularity of human corneal flaps prepared by femtosecond laser technology. Vossmerbaeumer U, Jonas JB. 24(6):645-648. [Report]
- Retinal nerve fiber layer measurements before and after photorefractive keratectomy. Iester M, De Feo F, Bricola G, Papadia M, Venturino G, Traverso CE, Calabria G. 24(6):639-644.
- Seven-year follow-up of LASIK for moderate to severe myopia. Liu Z, Li Y, Cheng Z, Zhou F, Jiang H, Li J. 24(9):935-940.

- Short chain aliphatic  $\beta$ -nitro alcohols for corneoscleral cross-linking: corneal endothelial toxicity studies. Paik DC, Wen Q, Braunstein RE, Trokel SL. 24(7):S741-S747. [Proceedings]
- Spontaneous bilateral, recurrent, late-onset diffuse lamellar keratitis after LASIK in a patient with Cogan's syndrome. Javaloy J, Barrera C, Muñoz G, Perez-Santonja JJ, Vidal MT, Alió JL. 24(5):548-550. [Report]
- Staged intrastromal delivery of riboflavin with UVA cross-linking in advanced bullous keratopathy: laboratory investigation and first clinical case. Krueger RR, Ramos-Esteban JC, Kanellopoulos AJ. 24(7):S730-S736. [Proceedings]
- Subconjunctival gas bubble formation during LASIK flap creation using femtosecond laser. Ide T, Kymionis GD, Goldman DA, Yoo SH, O'Brien TP. 24(8):850-851. [Report]
- Synergistic effect of ethanol and mitomycin C on corneal stroma. Netto MV, Barreto J Jr, Santo R, Bechara S, Kara-Jose N, Wilson SE. 24(6):626-632.
- The advanced human eye model (AHEM): a personal binocular eye modeling system inclusive of refraction, diffraction, and scatter. Donnelly W III. 24(9):976-983. [Proceedings]
- The effect of cyclosporine A (Restasis) on recovery of visual acuity following LASIK. Ursea R, Purcell TL, Tan BU, Nalgirkar A, Lovaton ME, Ehrenhaus MP, Schanzlin DJ. 24(5):473-476.
- The Visian myopic implantable collamer lens does not significantly affect axial length measurement with the IOLMaster. Sanders DR, Bernitsky DA, Harton PJ Jr, Rivera RP. 24(9):957-959. [Report]
- Topographic changes after hyperopic LASIK with the SCHWIND ESIRIS laser platform. de Ortueta D, Arba-Mosquera S, Baatz H. 24(2):137-144.
- Topography-guided custom ablation for irregular corneal astigmatism using the NIDEK NAVEX laser system. Yoshida Y, Nakamura T, Hara S, Yoshida N, Kojima T, Ichikawa K. 24(1):24-32.
- Tracking difficulties after femtosecond laser flap creation with the LADARVision excimer laser system. Ramos-Esteban JC, Bamba S, Krueger RR. 24(9):953-956. [Report]
- Transmission electron microscopy study of corneal epithelial flaps following removal using mechanical scraping, alcohol, and epikeratome techniques. Choi CY, Kim JY, Kim MJ, Tchah H. 24(7):667-670.
- Treatment of keratoconus by topography-guided customized photorefractive keratectomy: two-year follow-up study. Cennamo G, Intravaja A, Boccuzzi D, Marotta G, Cennamo G. 24(2):145-149.
- Tricyclic antidepressants: potential therapeutic alternatives for treatment of dry eye symptoms after LASIK. Ghaffariyeh A, Chamacham T. 24(8):770-771. [Letter]
- Tricyclic antidepressants: potential therapeutic alternatives for treatment of dry eye symptoms after LASIK. Tuisku IS, Tervo TM, Belmonte C. 24(8):771-772. [Reply to Letter]
- Troutman award for editorial excellence—16 years of inspired contributions. Waring GO III. 24(9):863-864. [Editorial]
- Undetected development of glaucoma after radial keratotomy. Scheuerle AF, Martin M, Voelcker HE, Auffarth G. 24(1):51-54. [Report]
- Urrets-Zavalía syndrome following iris-claw phakic intraocular lens implantation. Park SH, Kim SY, Kim HI, Yang SW. 24(9):959-961. [Report]
- Very high-frequency digital ultrasound measurement of the LASIK flap thickness profile using the IntraLase femtosecond laser and M2 and Carriazo-Pendular microkeratomes. Alió JL, Piñero DP. 24(1):12-23.
- Visual and refractive outcomes of LASIK with the SCHWIND ESIRIS and WaveLight ALLEGRETTO WAVE Eye-Q excimer lasers: a prospective, contralateral study. Mearza AA, Muhtaseb M, Aslanides IM. 24(9):885-890.
- Visual function and change in quality of life after bilateral refractive lens exchange with the ReSTOR multifocal intraocular lens. Blaylock JF, Si Z, Aitchison S, Prescott C. 24(3):265-273.
- Visual function with bilateral implantation of monofocal and multifocal intraocular lenses: a prospective, randomized, controlled clinical trial. Palmer AM, Faina PG, Albelda AE, Serrano MC, Saad DN, Cespedes MC. 24(3):257-264.
- Visual results following implantation of a refractive multifocal IOL in one eye and a diffractive multifocal IOL in the contralateral eye. Goes FJ. 24(3):300-305. [New Technology]
- Visualization of the retinal image in an eye model with spherical and aspheric, diffractive, and refractive multifocal intraocular lenses. Terwee T, Weeber H, van der Mooren M, Piers P. 24(3):223-232.
- Wavefront measurements of diffractive and refractive multifocal intraocular lenses in an artificial eye. Campbell CE. 24(3):308-311. [Report]
- Wavefront optimized versus custom-Q treatments in surface ablation for myopic astigmatism with the WaveLight ALLEGRETTO laser. Stojanovic A, Wang L, Jankov MR, Nitter TA, Wang Q. 24(8):779-789.
- Wavefront-guided custom ablation for myopia using the NIDEK NAVEX laser system. Venter J. 24(5):487-493.
- Wavefront-guided excimer laser ablation using photorefractive keratectomy and sub-Bowman's keratomileusis: a contralateral eye study. Durrie DS, Slade SG, Marshall J. 24(1):S77-S84. [Proceedings]
- Wavefront-guided excimer laser vision correction after multifocal IOL implantation. Jendritza BB, Knorz MC, Morton S. 24(3):274-279.
- Wavefront-optimized versus wavefront-guided LASIK for myopic astigmatism with the ALLEGRETTO WAVE: three-month results of a prospective FDA trial. Stonecipher KG, Kezirian GM. 24(4):S424-S430.
- What about LASEK? Camellin M. 24(5):462. [Letter]
- Where is the optimum far-point for a presbyopic eye? Thibos LN. 24(9):970-975. [Proceedings]